

67  
Cutaneous Sympathy

March 10

Jonas Underwood

admitted March 24<sup>th</sup> 1821

Coos Bay, Ore.

1901 May 1st to 1901



An Essay  
On Cutaneous Sympathy  
By Samuel Underwood M.D.

During  
the progre  
junction  
but, it is not  
skin it is  
to which, in  
its sensibility  
with the  
it. Previous  
actions and  
the instrument  
the medicine  
will be pro  
anatomical  
relation  
of binding  
parts, of acc  
matter, and  
between the  
It is divided



During the revolutions which have marked  
the progress of medical science, the secretory  
function of the skin has not been overlooked;  
but, it is only until within a few years, that the  
skin itself has received that degree of attention,  
to which its minute and complex structure,  
its sensibility and intimate sympathy  
with the various internal organs, entitles  
it. Previously to considering the associated  
actions maintained between the skin and  
the internal organs, and their connexion thro'  
the medium of the mucous membrane, it  
will be proper to present a brief view of the  
anatomical structure of the skin. This appa-  
rently simple envelope, serves the several purposes  
of binding together, and of protecting the internal  
parts, of excreting a large quantity of perspiratory  
matter, and of establishing an intimate relation  
between the living system and external substances.  
It is divided by anatomists into three distinct layers,

the per  
der, is the  
asolar dis  
with num  
surface, it  
watering  
free, wher  
quest co  
and the a  
water it b  
fibres, an  
Examine  
penetrate  
the pump  
the hairs  
tissue of  
myriads  
close cap  
compact  
network over

the principle of which, constituting the chief bulk of the  
 skin, is the *cutis vera*. This is a compact and strong  
 areolar tissue, composed of a firm fibrous substance,  
 with numerous vacuities or intervals. On its external  
 surface, it is close and compact resembling the smooth  
 continuity of a membrane, but on its opposite sur-  
 face, where the fibrous portion is blended with sub-  
 jacent cellular substance, the texture is more loose,  
 and the areolae are larger. When immersed in  
 water it becomes softer, by a separation of its  
 fibres, and its intervals are rendered more distinct.  
 Examined in this state, the areolae are found to  
 penetrate its whole thickness, and to serve for  
 the purpose of transmitting to the surface  
 the hairs, exhalants and absorbents. The areolar  
 tissue of the *cutis vera* is permeated by countless  
 myriads of arterial and venous ramifications,  
 whose capillary divisions occupy the external or  
 compact surface of this part, and form a vascular  
 network over the whole body. This is rendered obvious in

the glow of  
position of  
Saxanthus  
exchange  
modern of  
manifested  
in those  
invisible  
nearly equal  
mucus of  
larger ram  
hillside. A  
is the thin  
blister. See  
to the subg.  
laminar.  
Roodness of  
hairs, by the  
by the exha  
sily in some

the glow of aruse and the flush of shame; in the excitement of fever and in the eruptive stage of Exanthematous diseases; when this part becomes surcharged with blood. It may also be proved post mortem by the injection of coloured fluids. These vascular ramifications are found to be particularly numerous in those parts, which possess most exquisite sensibility. The absorbents of the skin are nearly equal in number to its bloodvessels, and mucous nerves enter it in all its parts, distributing their larger ramifications in situations occupied by the papillae. The cuticle, the anterior layer of the integument, is the thin transparent pellicle, which is raised by a blister. In its natural state, it adheres almost inseparably to the subjacent parts. It presents no traces of fibres, laminae or cells, and, in it, we can discover neither bloodvessels, absorbents nor nerves. It is perforated by hairs, by the excretory tubes of the cutaneous follicles, by the exhalant mouths of the capillary vessels, and, probably in some parts, by absorbent orifices. It is insensible

and sub-  
stitutions, and  
inorganic  
would have  
placed as  
and delicate  
medium of  
impression  
internation  
various a  
contrast of  
impart to  
The roman  
is a delicate  
slightly de  
rium. It h  
of its texture  
is produced  
and other  
rat of all the

and supposed incapable of performing any vital actions, and has been, emphatically, denominated, inorganic and avascular. The external covering of the leg would, hence, appear to be but a dead membrane, placed as a protection to the finely organised and delicate parts beneath, and as an insensible medium for the communication of external impressions; for we find if this part be thickened preternaturally, that sensation is in a great measure destroyed, and, if it be removed, the contact of bodies gives pain and does not impart the appropriate impression of touch. The remaining portion of the skin may be regarded as a delicate stratum interposed between the parts already described, and is denominated cutis mucosa. It has derived its name from the softness of its texture and its netlike appearance, which is produced by the perforations of hairs, papillae and other parts coming to the surface. It is the seat of all the variety of colour which has been observed in

*[Faint, illegible handwriting, likely bleed-through from the reverse side of the page.]*

the human  
and brain  
of the body  
essentially  
one great  
external  
it is observed  
constant  
and, are  
absorbed  
the vascular  
in all the  
the symphysis  
external  
between the  
Culano  
marked, that  
almost  
some of our



the human species. The skin considered as one membrane may be regarded as the sensitive limit of the body. Placed at the extremity of its organs, and incessantly exposed to external influence, it forms one great connexion between animal existence and external objects. From this brief anatomical view, it is obvious, that the skin is adapted to hold important relations with various other parts of the system. In it, are represented the nerves, arteries, veins, and absorbents, especially the capillary portion of the vascular system, which is so essentially concerned in all the vital functions. I shall next point out the sympathies of this part with some of the internal organs. The first, which materially presents itself to our notice, is that, which exists between the skin and the lungs, denominated Cutaneo Pulmonary sympathy. This is so strongly marked, that to insist upon it, would seem almost superfluous; but as it is the source of some of our most formidable diseases, I shall

consider  
agent, of  
of discon  
by moie  
Here and  
ent port  
United St  
the town  
portion,  
vicissitud  
all the q  
dominanc  
great ex  
vicissitud  
and the p  
do not ob  
consider  
and a pr  
effusions  
or less freq

consider it somewhat in detail. The principle agent, operating on the skin in the production of disease, is atmospherical vicissitudes, aided by moisture. A reference to the state of the atmosphere, and to the diseases peculiar to the different portions of the Atlantic division of the United States, will incontestably establish the truth of this position. In the eastern portion, where we have great and sudden vicissitudes, with much moisture, we find all the forms of pulmonary disease predominating and prevailing to a very great extent. In the middle portion, where these vicissitudes are still sudden and considerable, and the moisture is also great, but where they do not obtain to the same extent, there is a considerable decrease of pulmonary disease and a proportional increase of rheumatic affections. In the southern states, where these changes are less frequent, and the country is, in a great measure,

allusion  
nature, i  
to. Sorta  
bilious  
have b  
Pulson  
of the  
in the  
of the  
from  
arrest  
I have  
below the  
surface  
ed to b  
a torpe  
the less  
transm  
fucose  
in all b

alluvial and abounding in marshes and stagnant waters, we find these forms of disease giving place to Intermitting and all the various forms of bilious disease. These portions of the United States have been emphatically denominated the Pulmonic, Rheumatic and Bilious sections of the Atlantic states. The effects of temperature on the Pulmonic system through the medium of the skin, independently of the proofs derived from daily observation, may be illustrated by direct experiment. It is observed by Mr. James Johnson that on immersing the body in water below the temperature of the skin, the vessels on the surface are struck torpid and the blood is determined to the interior. At this moment a synspasm is torpor takes place in the capillary vessels of the lungs, so that the blood is with difficulty transmitted thro' them, occasioning that Dyspnoea, or panting for breath, which we observe in all but more particularly in delicate persons

*[Faint, illegible handwriting across the page]*

at  
1000  
is 1000  
new  
10/100  
But  
more  
it was  
pass  
1000  
over  
circ  
and  
are  
accor  
The  
exce  
the fa  
1000  
the co

at the moment of immersion. But here reaction soon takes place. The balance of the circulation is restored and the functions of the skin are renewed with increased activity, succeeded by exhilaration of spirits and renovated energy. But should there have been previously to immersion exercise sufficient to induce fatigue or excess of perspiration to weaken the extreme vessels on the surface, then the torpor of the extreme vessels of the skin cannot be properly overcome by reaction, the balance of the circulation is not completely restored, and the lungs or other internal organs are injured, attended with more or less fever according to the force of the operating cause. The sympathy, I am contending for, exercises receives additional support from the fact, that no means have yet been discovered, so invariably successfully in the cure of chronic Pulmonary disease, as

the apper  
of the body  
even flou  
skin, and  
then any  
cases a h  
proof, en  
drawn  
which do  
ting up  
through  
from the  
damps of  
have bee  
on pale  
education  
sympath  
associat  
and the



the application of nonconductors to the surface of the body. In a great number of cases, I have seen flannel prescribed to be worn next the skin and in numerous slight cases, have found <sup>it</sup> amply sufficient for the cure, and, in all cases, a highly useful auxiliary. Another proof, in support of this sympathy, may be drawn from the enormous waste of life, which diurnal vicissitudes occasion, by operating upon the delicate lungs of females through the medium of the skin. This results from their frequent exposure to the chilly damps of night, after the perspiratory vessels have been overexcited in crowded rooms, or fatigue has been induced by the seductive exertions of the dance.

The Cutaneous Gastric sympathy, as its name imports, is the association of action between the skin and the stomach. This sympathy

has been  
by Cullen  
causes a  
that pro  
ficients  
between  
the body  
appears  
state of  
appetite  
applied  
it does  
dissected  
means of  
fore allo  
vomitory  
to this ca  
adduces  
his descrip  
and only

has been particularly observed and described by Cullen, in his exposition of the proximate causes and phenomena of fever. He remarks, that from many circumstances it is sufficiently certain, that there is a consent between the stomach and the surface of the body; that this consent particularly appears, from the connexion between the state of perspiration and the state of the appetite in healthy persons; that cold, applied to the surface of the body where it does not stop perspiration, proves a stimulus to the surface, and is a powerful means of exciting appetite. Dr. Cullen therefore attributes the anorexia, nausea, and vomiting, which occur in febrile affections to this cause; and in support of this doctrine adduces the fact related by Sydenham in his description of Plague, in which, vomiting could only be allayed by exciting the surface, so as to

induce  
illustrated  
in the sto  
ation of  
few who h  
Delicate  
shoes are  
although  
in food a  
more sub  
all that  
arise. fo  
so strong  
with the  
died from  
called the  
But the im  
make upon  
their rece  
most essen

induce perspiration. This sympathy is further illustrated by the frequent occurrence of pain in the stomach and indigestion from the application of cold and moisture to the feet. There are few who have not suffered more or less from this cause. Delicate females with tight dress and thin shoes are particularly obnoxious to it. Women, although they are infinitely more temperate in food and drink than men, yet they are more subject to Cardialgia, Flatulence and all that train of distressing symptoms, which arise from a disordered state of the stomach. So strongly impressed was the late Dr. Rush with the belief, that numerous diseases originated from this source, that he emphatically called the feet "the high road to disease." But the impression which some medicines evidently make upon the skin, in a very short time after their reception into the stomach, affords the most remarkable proof of this connexion;

and  
the mea  
pitrous  
conven  
and  
the inter  
which  
-inated  
is severely  
intention  
al impor  
-tion and  
frequen  
a highly  
quently  
tudes or  
-sequent  
forms of  
places,  
of count

and here I speak of those, which act not through the medium of the circulation. e.g. I have seen nitrous powders, in a quarter of an hour convert a hot, parched skin into a perspiring one. The next association of the skin with the internal organs, which I shall notice is that, which exists between it and the intestines denominated Cutaneo-Intestinal sympathy. There is scarcely any cause of functional disorder in the intestines, so common as that resulting from external impressions on the skin. Suppressed perspiration and cold or moisture applied to the feet, frequently induce Diarrhoea & Colic, and a highly dangerous state of Enteritis is frequently brought on by atmospherical vicissitudes or cold applied to the skin during or subsequent to a state of perspiration. The milder forms of Dysentery are met with in most places, but in elevated ridges of low districts of country, or in which marsh effluvia and

*[Faint, illegible handwriting on the left page, likely bleed-through from the reverse side.]*

...dines per  
disease  
malign  
and ala  
which is  
they wido  
conceder  
colica  
of the inst  
legs and  
put upon  
symptom  
between  
chiefly a  
of the vis  
a state  
should  
for we p  
other dis  
are irritat



atmospherical vicissitudes are combined, this disease frequently assumes an obstinate and malignant form, which renders it as mortal and alarming, as those fatal epidemics, which savage our large cities. This sympathy will be rendered more apparent, from considering the great relief afforded in Colica tritonum and other spasmodic diseases of the intestines, by decubating cold water on the legs and thighs or from standing with bare feet upon cold pavements. The cutaneous renal sympathy is that connection, which exists between the skin and the kidneys, and is chiefly deserving of consideration on account of the vicarious actions of these organs during a state of health. This sympathy, however, should not be lost sight of in disease, for we find, that calculous affections, as also other diseases, in which the kidneys are concerned are irritated and increased by exposure to cold, and

that of  
warm  
shall not  
and live  
by. The  
early de  
work on  
was by a  
distinct  
are most  
mark the  
between the  
the liver.  
climates, a  
regions con  
invention  
perpetuate  
ing dimin  
complicated

that few remedies have been so effectual as the warm bath. The last sympathy, which I shall notice is that, which exists between the skin and liver, called the Cutano Hepatic sympathy. This was distinctly noticed and particularly described by Mr. Johnson in his valuable work on the diseases of tropical climates. It was, by attentively observing diseases in those districts of country, where hepatic affections are most abundant, that he was led to remark the very intimate relation, which exists between the functions of the skin and those of the liver. It is a well known fact, that, in tropical climates, and in the hot season of more temperate regions, when perspiration is abundant, biliary secretion is also in excess, and that checked perspiration is always attended with a corresponding diminution of biliary secretion —

— A high degree of temperature by augmenting the

citizenship  
usult by  
and hence  
tropical by  
find the  
hot climatic  
temperate  
the night  
ably acc  
Such are  
of the who  
part of the  
time, which  
have not  
production  
beyond the  
national  
indiscrimi  
ity. It has  
associated

cutaneous and biliary secretion debilitates the  
 vessels, by which these processes are carried on,  
 and hence they are the more easily reordered  
 torpid by the application of cold. Accordingly we  
 find Hepatitis, Cholera Morbus &c. to prevail in  
 hot climates, and during the hot season in more  
 temperate regions, and also to occur during the  
 the night season, when the atmosphere is conside-  
 rably cooler than that of the preceding day.  
 Such are some of the surprising sympathies  
 of the skin with the internal organs. The  
 fact of their existence and the important indica-  
 tions, which they point out in the cure of diseases  
 have not been overlooked, but the mode of their  
 production has been deemed inexplicable and  
 beyond the ken of human research. This I ap-  
 prehend has arisen, in part, from the vague and  
 indiscriminate application of the term Sympa-  
 thy. It has been unable to denote, not only, the  
 associated action of parts, between which a

hood  
can be  
no such  
sympat  
there is  
us with  
than is  
vaccin  
in their  
skin, an  
organ. &  
the skin  
the body  
demon  
of the ho  
-nally.  
memb  
surface  
and too  
and to

direct connexion, in anatomical structure,  
 can be traced, but also of those parts where  
 no such connexion exists. Upon reviewing these  
 sympathies of the skin with the internal organs,  
 there is one circumstance, which must impress  
 us with the idea of a connexion less mysterious  
 than is usually imagined. All the different  
 viscera, which have been just stated to respond  
 in their actions to the impressions made on the  
 skin, are lined by a direct extension from this  
 organ. The mucous membrane passes from  
 the skin through all the natural apertures of  
 the body to become, as Bichat has sufficiently  
 demonstrated, the internal cutaneous system  
 of the hollow viscera, which communicate exter-  
 nally. The lungs are supplied with this lining  
 membrane, which after covering the internal  
 surface of the fauces, passes down into the larynx  
 and trachea, and, of course, into the bronchiae,  
 and thus supplies their whole internal surface.

The liver  
for the m  
tissue of  
solidated  
seri bilina  
brane, fas  
and tie b  
membran  
and funic  
the parts, p  
preserves  
animal on  
the mesop  
nigrum, w  
of the exte  
of colour  
laminar  
gus. As  
processes  
the skin, s



7  
The liver is also lined in the same manner;  
for the mucous coat of the duodenum is con-  
tinued up through the ductus communis  
choledochus both into the gall bladder and the  
pore biliaris. In like manner the mucous mem-  
brane passes thro' the ureters into the infundibulum  
and tubuli uriniferi of the kidneys. This  
membrane, though modified in its structure  
and functions, according to the necessities of  
the parts, to which it is appropriated, still  
preserves, in its remotest passages, the  
general characteristics of the skin, with  
the exception of the peculiar pigmentum  
nigrum, which imbedded in the rete mucosum  
of the external skin, produces the varieties  
of colour thro' the translucent cuticle. The  
laminae of both systems are exactly analo-  
gous. As the cuticle sends down its vaginal  
processes through the glandular follicles of  
the skin, so the epithelium of the mucous tissues

gives a co  
the hollow  
prominate  
of the cutis  
the corioepi  
membrane.  
we found  
them by its  
external bea  
tions of the  
of the intes  
of solid an  
to the perspe  
surface, and  
innervated  
mucous fl  
the intima  
a general  
will know  
each syste

gives a corresponding lining through all the hollow passages. The vascular tissue, demonstrated by Bagnard upon the surface of the uctis, is asserted by Richat to exist between the corresponding laminae of the whole mucous membrane. In both tissues the vessels and nerves are found to ramify in exact similarity. As the skin by its nervous papillae takes cognizance of external bodies in touch, so the analogous terminations of the nervous fibrils in the different portions of the interior membrane convey the impressions of sapid and odouriferous particles to the mind. As the perspiratory fluid is secreted on the external surface and performs the office of maintaining the associated functions in exactest harmony, so the mucous fluids secreted on their surfaces, keep up the similar motions peculiar to each organ. It is a general law of the animal oeconomy, too well known to need much elucidation, that each system of parts is naturally disposed

to what  
must. The  
system  
takes in  
some de  
vacuatio  
this ind  
and sy  
the alim  
most es  
introduc  
vacuati  
they can  
imitatio  
invested  
the stor  
Indolence  
bodies in  
duce sev  
many a

to act in the same state of energy and excitement. Thus an irritation in any part of the arterial system, though at first only the neighboring tubes may be excited, eventually produces the same degree of action throughout the whole circulation. It is the struggle to accomplish this end, which creates the inflammatory and sympathetic fever of wounds &c. In the alimentary canal, this law is however most remarkably illustrated. Some purgatives introduced into the stomach excite a general evacuation of the large intestines long before they can travel beyond the pylorus. The irritation in the fauces or oesophagus excites inverted motions for its discharge; and often the stomach joins in the same action.

Indolent hemispherical tumours, and hard bodies in the rectum have been known to produce severe intestinal derangement, and many obstinate Dyspepsias have depended

upon the  
when we  
into the  
the inter  
organs, a  
external  
atomistic  
shown to  
erve tis  
An imp  
surface  
there, th  
monia  
junction  
supposed  
the same  
consider  
which I  
all the

upon the existence of fistulas in ano. Nor when we recollect, that the skin extends itself into the mucous membrane, and thus forms the internal secreting surface of all those organs, which communicate with it by the external apertures, it will not appear more astonishing that these viscera, which I have shown to be sympathetically <sup>connected</sup> with this extensive tissue, should show the same disposition. An impression made upon one part of its surface and producing a consequent action there, the whole organ wents itself to harmonize in the same motions: and these functions, which at first sight might be supposed widely diverse, are operating under the same excitement, and may surely be considered as subject to the general law, which I have maintained as influencing all the other systems.

J. Underwood. No 266 Archd.

Dr. Chapman

My son  
er

The sym

The